

BACH2 mediates negative selection and p53-dependent tumor suppression at the pre-B cell receptor checkpoint.

Journal: Nat Med

Publication Year: 2013

Authors: Srividya Swaminathan, Chuanxin Huang, Huimin Geng, Zhengshan Chen, Richard Harvey, Huining Kang, Carina Ng, Bjorn Titz, Christian Hurtz, Mohammed Firas Sadiyah, Daniel Nowak, Gabriela B Thoenissen, Vikki Rand, Thomas G Graeber, H Phillip Koeffler, William L Carroll, Cheryl L Willman, Andrew G Hall, Kazuhiko Igarashi, Ari Melnick, Markus Muschen

PubMed link: 23852341

Funding Grants: Dual targeting of tyrosine kinase and BCL6 signaling for leukemia stem cell eradication

Public Summary:

The B cell-specific transcription factor BACH2 is required for affinity maturation of B cells. Here we show that Bach2-mediated activation of p53 is required for stringent elimination of pre-B cells that failed to productively rearrange immunoglobulin VH-DJH gene segments. After productive VH-DJH gene rearrangement, pre-B cell receptor signaling ends BACH2-mediated negative selection through B cell lymphoma 6 (BCL6)-mediated repression of p53. In patients with pre-B acute lymphoblastic leukemia, the BACH2-mediated checkpoint control is compromised by deletions, rare somatic mutations and loss of its upstream activator, PAX5. Low levels of BACH2 expression in these patients represent a strong independent predictor of poor clinical outcome. In this study, we demonstrate that Bach2(+/-) pre-B cells resist leukemic transformation by Myc through Bach2-dependent upregulation of p53 and do not initiate fatal leukemia in transplant-recipient mice. Chromatin immunoprecipitation sequencing and gene expression analyses carried out by us revealed that BACH2 competes with BCL6 for promoter binding and reverses BCL6-mediated repression of p53 and other cell cycle checkpoint-control genes. These findings identify BACH2 as a crucial mediator of negative selection at the pre-B cell receptor checkpoint and a safeguard against leukemogenesis.

Scientific Abstract:

The B cell-specific transcription factor BACH2 is required for affinity maturation of B cells. Here we show that Bach2-mediated activation of p53 is required for stringent elimination of pre-B cells that failed to productively rearrange immunoglobulin VH-DJH gene segments. After productive VH-DJH gene rearrangement, pre-B cell receptor signaling ends BACH2-mediated negative selection through B cell lymphoma 6 (BCL6)-mediated repression of p53. In patients with pre-B acute lymphoblastic leukemia, the BACH2-mediated checkpoint control is compromised by deletions, rare somatic mutations and loss of its upstream activator, PAX5. Low levels of BACH2 expression in these patients represent a strong independent predictor of poor clinical outcome. In this study, we demonstrate that Bach2(+/-) pre-B cells resist leukemic transformation by Myc through Bach2-dependent upregulation of p53 and do not initiate fatal leukemia in transplant-recipient mice. Chromatin immunoprecipitation sequencing and gene expression analyses carried out by us revealed that BACH2 competes with BCL6 for promoter binding and reverses BCL6-mediated repression of p53 and other cell cycle checkpoint-control genes. These findings identify BACH2 as a crucial mediator of negative selection at the pre-B cell receptor checkpoint and a safeguard against leukemogenesis.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/bach2-mediates-negative-selection-and-p53-dependent-tumor-suppression-pre-b>